

Sensitivity to the $X(3872)$ total width at the Belle II experiment


Hikari Hirata (Nagoya University)
for the Belle II collaboration

FPCP 2019

University of Victoria, May 9th 2019

X(3872)


- One candidate of exotic hadrons.
 - It was discovered in **B decay**.




$c\bar{c}$ component is included.

ρ meson component ($I = 1$) is dominant
 $\rightarrow u\bar{u}/d\bar{d}$ is also included.

Phys. Rev. Lett. 91, 262001




BELLE-CONF-0541



Cannot be interpreted
 as pure charmonium.

X(3872)


- One candidate of exotic hadrons.
 - It was discovered in **B decay**.



$c\bar{c}$ component is included.

ρ meson component ($l=1$) is dominant
 $\rightarrow u\bar{u}/d\bar{d}$ is also included.

- Various decay modes are observed.
- $M_{X(3872)} = 3871.69 \pm 0.17 \text{ MeV}/c^2$
 \rightarrow Consistent with $D^0\bar{D}^{*0}$ threshold within the error.
- $J^{PC} = 1^{++}$




Branching ratio

\rightarrow Support interpretation as pure $D\bar{D}^*$ molecule.

Counter evidence of pure molecule model

Cross section of prompt X(3872)
in pp-collision (LHC experiment)



- Should be suppressed for a molecule.
→ Consistent with cross section for pure $c\bar{c}$ state ($\chi_{c1}(2P)$).
→ $D\bar{D}^* - c\bar{c}$ mixture state?
- Need further information about production and decay.
→ Current X(3872) total width:

$$\Gamma_{\text{tot}} < 1.2 \text{ MeV}$$



We aim to measure **significant X(3872) total width.**

→ Provides us **partial widths for each decay mode.**

$$\Gamma(X(3872) \rightarrow f) = \frac{\text{Br}(B^\pm \rightarrow K^\pm X(3872)) \times \text{Br}(X(3872) \rightarrow f)}{\text{Br}(B^\pm \rightarrow K^\pm X(3872))} \times \Gamma_{\text{tot}}$$

Belle II experiment

- An electron-positron collider B factory
 $\sqrt{s} = 10.58 \text{ GeV}$



Belle II detector

Central Drift chamber

Vertex Detector

2 layers Si pixels
+4 layers strips

EM calorimeter

Cs(Tl) crystal

Particle Identification

[Forward] Aerogel-RICH
[Barrel] Time of Propagation

Solenoid

1.5 T

K_L and μ detector

Capable of detecting π^\pm , K^\pm , p , e , μ and γ for wide momentum region.

- Goal of integrated luminosity:
 $50 \text{ ab}^{-1} = \sim 5.5 \times 10^{10} \text{ B meson pairs}$ ($\times 50$ that of Belle).

Strategy for improvement of sensitivity

- Previous study (Phys. Rev. D 84, 052004 (2011))
 - Use $X(3872) \rightarrow J/\psi\pi^+\pi^-$ mode.
 - Fit signal component of mass spectrum with Breit-Wigner convoluted with mass resolution.


$$\Gamma_{\text{tot}} < 1.2 \text{ MeV (90\% C.L.)}$$

$$<$$

$$\text{Mass resolution } 1.86 \pm 0.01 \text{ MeV}/c^2$$

“Improvement of mass resolution is essential”

- ★ In general, decay mode with small Q-value has good mass resolution.



- ▶ However, the signal yield is too low to measure total width so far.


Strategy for improvement of sensitivity

- Previous study (Phys. Rev. D 84, 052004 (2011))
 - Use $X(3872) \rightarrow J/\psi\pi^+\pi^-$ mode.

Belle II is suitable for this measurement thanks to huge data sample.

In this presentation, analysis overview and sensitivity of total width with simulation will be shown.


resolution.



- ▶ However, the signal yield is too low to measure total width so far.

Reconstruction and selection

Decay chain:



- D^0 mode

$$\begin{cases} D^0 \rightarrow K^+ \pi^- \\ D^0 \rightarrow K^+ \pi^- \pi^0 \\ D^0 \rightarrow K^+ \pi^- \pi^- \pi^+ \end{cases}$$

~ 26%
 D^0 modes

- π^0 mode

$$\pi^0 \rightarrow \gamma\gamma$$

~ 99%
 π^0 modes


Event selection

- For final state particle, PID, tracking (K^\pm , π^\pm) and cluster information (γ).
- For D^0 and π^0 mesons, signal regions are selected, and mass-constrained fits are used.
- For B mesons, beam-energy-constrained mass and CMS energy difference.
- In order to reduce multiplicity of B candidates, best candidate selection is performed by selecting a candidate with minimum χ_{BCS}^2 .

$$\chi_{BCS}^2 = \left(\frac{\Delta M_{D^0}}{\sigma M_{D^0}} \right)^2 + \left(\frac{\Delta M_{\bar{D}^0}}{\sigma M_{\bar{D}^0}} \right)^2 + \left(\frac{\Delta M_{\pi^0}}{\sigma M_{\pi^0}} \right)^2 + \left(\frac{E_B - E_{\text{beam}}}{\sigma_{(E_B - E_{\text{beam}})}} \right)^2$$

$D^0\bar{D}^0\pi^0$ invariant mass

Mass spectrum after reconstruction and selection



- **Correct** reconstructed of signal
- **Incorrect** reconstruction of signal
- BG from general $B\bar{B}/q\bar{q}$ events

- Mass resolution: 684 ± 8 keV
- Signal yield with 1 ab^{-1} :
 $64.5 \pm \underline{23.9}$

Comes from large error of
 $\text{Br}(B^\pm \rightarrow K^\pm X(3872))$
 $\times \text{Br}(X(3872) \rightarrow D^0\bar{D}^0\pi^0)$



→ Compared with previous study ($J/\psi\pi^+\pi^-$ mode),
 Signal yields is around half, but the mass resolution is 3 times
 better.

$D^0\bar{D}^0\pi^0$ mode provides a sample with the good mass resolution.

Total width extraction


- Total width is extracted by fitting the mass spectrum.
- Check if the total width is obtained by the fit correctly.
 - Linear relation between the total width generated in the simulation and that obtained by fit.

“Confirmed that there is no bias in the fit.”



Sensitivity to total width of X(3872)

- Sensitivity is estimated with toy-MC samples.



- With the full data sample of Belle II (50 ab^{-1}), total width with values up to
[90% C.L.] ~ 180 keV
[3 σ significance] ~ 280 keV
[5 σ significant] ~ 570 keV
 can be measured.

- Belle II is capable of measuring total widths.
- Next, we need detailed study of the effect of mass resolution and background (shape and beam background)
 → Suppress systematics.

Conclusion

- In order to derive $X(3872)$ partial widths for each decay mode, we aim at significant measurement of $X(3872)$ total width.
- Sensitivity to the total width at Belle II is estimated by simulation
 - We used a sample with good mass resolution, $D^0\bar{D}^0\pi^0$ decay mode.
→ There is no bias in total width extraction.
 - With the full data sample of Belle II, it is possible to measure total width with $3\sigma(5\sigma)$ significance with values up to 280(570) keV.
- Next, we need detailed study of effect of mass resolution and background (shape and beam background).
- We also plan to measure precise mass spectrum to test possibility of cusp.


Stay tuned!

Thank you for your attention.

Partial width for $X(3872) \rightarrow J/\psi\gamma$

$$\Gamma(X(3872) \rightarrow J/\psi\gamma) = \Gamma_{tot} \times \underbrace{BR(X(3872) \rightarrow J/\psi\gamma)}_{< \sim 1\%} < 0.01 \times \Gamma_{tot}$$

If Γ_{tot} upper limit = 180 keV, $\Gamma(X(3872) \rightarrow J/\psi\gamma) < 18$ keV



[1] S. Dubnicka, et. al., Phys. Rev. D **81**, 114007 (2010)


[2] F. Aceti, et. al., Phys. Rev. D **86**, 113007 (2012)

[3] Y. Dong, et. al, J. Phys. G: Null.Part. Pays. 38, 015001 (2011)

Partial width for $X(3872) \rightarrow J/\psi\gamma$

$$\Gamma(X(3872) \rightarrow J/\psi\gamma) = \Gamma_{tot} \times \underbrace{BR(X(3872) \rightarrow J/\psi\gamma)}_{< \sim 1\%} < 0.01 \times \Gamma_{tot}$$

If Γ_{tot} upper limit = 180 keV, $\Gamma(X(3872) \rightarrow J/\psi\gamma) < 18$ keV




[1] S. Dubnicka, et. al., Phys. Rev. D **81**, 114007 (2010)

[2] F. Aceti, et. al., Phys. Rev. D **86**, 113007 (2012)

[3] Y. Dong, et. al, J. Phys. G: Null.Part. Pays. 38, 015001 (2011)


Br($B^\pm \rightarrow K^\pm X(3872)$) measurement



X_{cc} is reconstructed by missing mass

$$M_{X_{cc}}^2 = (P_{\text{beam}} - P_{B_{\text{tag}}} - P_{K^+})^2$$

$$\rightarrow \text{Br}(B^\pm \rightarrow K^\pm X(3872)) < 2.6 \times 10^{-4}$$




- With the full data sample of Belle II, it is possible to measure with 7σ significance (naive expectation).
- More realistic simulation is on going.

Statistical and systematic uncertainty

- Systematics $\sim +200/-660$ keV (Phys. Rev. D **96**, 074014 (2017))

* Naive expectation from previous study of $D^0\bar{D}^{*0}$ mode



- Statistical uncertainty


Summary table of total width sensitivity and statistical error obtained by fit

Integrated lumi. [ab^{-1}]	3σ significance [keV]	Statistical error [keV]
10	1000	± 550
20	580	± 140
30	440	± 150
40	360	± 90
50	280	± 80

- With $>20 \text{ ab}^{-1}$, systematic error become dominant \rightarrow Make it suppress

Relation between mass resolution and Q value

- Consider a decay mode of particle A, $A \rightarrow bcd\dots$
- Reconstructed mass M



$$M = \sqrt{\left(\sum_{i=bcd\dots} E_i\right)^2 - \left(\sum_i \vec{P}_i\right)^2}$$

→ Mass resolution σM can be derived as follows

$$\sigma M = \sqrt{\sum_i \left(\frac{\partial M}{\partial E_i}\right)^2 (\sigma E_i)^2 + \sum_i \left(\frac{\partial M}{\partial P_{x_i}}\right)^2 (\sigma P_{x_i})^2 + \dots}$$

$$\left\{ \begin{array}{l} \frac{\partial M}{\partial E_i} = \frac{E_i}{M} \\ \frac{\partial M}{\partial P_{x_i}} = \frac{P_{x_i}}{M} \\ \vdots \end{array} \right.$$

For the smaller Q-value mode, E , P_{x_i} , P_{y_i} , P_{z_i} are smaller.

→ Therefore, the mass resolution are small.