




# Prospects in spectroscopy with Belle II

Vishal Bhardwaj  
IISER Mohali  
*(for Belle II collaboration)*

14-19 July 2018

16<sup>TH</sup> CONFERENCE ON FLAVOR PHYSICS & CP VIOLATION


**FPCP 2018**



# Outline of the talk


- ❖ Motivation for spectroscopy
- ❖ Spectroscopy at  $B$  factories
- ❖ Belle to Belle II
- ❖ Prospects of charmonium spectroscopy in Belle II
- ❖ Bottomonium spectroscopy prospects
- ❖ Summary

# QCD : real particles are color singlet



Baryons are red-blue-green triplets  
 $\Lambda = usd$

Mesons are color-anticolor pairs




$\pi = \bar{u}d$

Other possible combinations of quarks and gluons : **eXoTiC**

*artistic illustration*


## Pentaquark

$S = +1$   
Baryon




## H di-Baryon

Tightly bound  
6 quark state




## Glueball

Color-singlet multi-gluon bound state




## Tetraquark

Tightly bound  
diquark &  
anti-diquark




## Molecule

loosely bound  
meson-  
antimeson  
"molecule"





$q\bar{q}$  -gluon hybrid  
mesons




- $q\bar{q}$  spectroscopy with heavy quark (mostly  $c$  or  $b$ ) are best place to study quark model.
- Simple two body system, non-relativistic and narrow (with OZI suppression),
- Further, one can search for exotics with them.


# Production of $q\bar{q}$ (-like) @ $B$ -factories



Reconstruct  $J/\psi$  and look at recoil mass






## Initial state radiation





Annihilation at smaller energy.

## Quarkonium decay/transitions



# $q\bar{q}$ (-like) states till now




- $1\frac{1}{2}$  decade has passed after the discovery of first  $c\bar{c}$ -like [  $X(3872)$  ] by the Belle collaboration.
- Plenty of states have been found.
- Several states found in one process (not easy to understand).
- States have non-zero charge, suggesting them to be tetraquark/molecule-like state.
- Instead of conventional spectroscopy, it is now *eXotiC spectroscopy*.
- However, the limited statistics always come as the evil limiting factor.



Belle II (with ability to accumulate 50 times\* more data in comparison to Belle) can play crucial role in understanding these states.

# Belle to Belle II




PROJECT	KEK/KEK	KEK	KEK	KEK
NAME	KEK/KEK	KEK/KEK	KEK/KEK	KEK/KEK
DATE	2011.03.25	2011.03.25	2011.03.25	2011.03.25
REV.	1	1	1	1
SCALE	1/10	1/10	1/10	1/10
FIG.	A	A	A	A

Belle II 6



# Belle to Belle II

Vertex detector  
 4 SVD layer  $\rightarrow$  2 layers  
 DEPFET + 4 layers DSSD  
 Expected resolution of  
 $\sim 25\mu\text{m}$  while in Belle  $\sim 50\mu\text{m}$




SYSTEM	REVISION	DATE	APPROVED BY	DATE
DESIGN	1/10	2008/08	Y. ITO	2008/08
DESIGNER	Y. ITO			
REVISION	1/10			
Belle II /				1/10

# Belle to Belle II

Vertex detector  
 4 SVD layer → 2 layers  
 DEPFET + 4 layers DSSD  
 Expected resolution of  
 ~25μm while in Belle ~50μm

CDC  
 Larger volume drift chamber,  
 smaller drift cell. Faster  
 electronics



SYSTEM	REVISION	DATE	APPROVED BY	DATE
DESIGN	1/10	2006/08/08	Y. ITOH	2006/08/08
DESIGNER	Y. ITOH	DATE	APPROVED BY	DATE
Y. ITOH	1/10	2006/08/08	Y. ITOH	2006/08/08

Page 8




# Belle to Belle II

**Vertex detector**  
 4 SVD layer  $\rightarrow$  2 layers  
 DEPFET + 4 layers DSSD  
 Expected resolution of  
 $\sim 25\mu\text{m}$  while in Belle  $\sim 50\mu\text{m}$

**CDC**  
 Larger volume drift chamber,  
 smaller drift cell. Faster  
 electronics

**PID**  
 More compact. Time of  
 Propagation (barrel) and prox.  
 foc. ARICH (Endcap) is used.  
 Provide better  $K/\pi$  separation.




# Belle to Belle II

**Vertex detector**  
 4 SVD layer → 2 layers  
 DEPFET + 4 layers DSSD  
 Expected resolution of  
 ~25μm while in Belle ~50μm

**CDC**  
 Larger volume drift chamber,  
 smaller drift cell. Faster  
 electronics

**PID**  
 More compact. Time of  
 Propagation (barrel) and prox.  
 foc. ARICH (Endcap) is used.  
 Provide better K/π separation.

**ECL**  
 Old crystals are used with  
 modified waveform sampling  
 electronics to reject pile-up  
 events.



# Belle to Belle II

## Vertex detector

4 SVD layer  $\rightarrow$  2 layers  
DEPFET + 4 layers DSSD  
Expected resolution of  
 $\sim 25\mu\text{m}$  while in Belle  $\sim 50\mu\text{m}$

## CDC

Larger volume drift chamber,  
smaller drift cell. Faster  
electronics

## PID


More compact. Time of  
Propagation (barrel) and prox.  
foc. ARICH (Endcap) is used.  
Provide better  $K/\pi$  separation.

## KLM

Resistive place counter  
(Barrel)  
Scintillation + WLSF + MSCP  
(endcap)

## ECL

Old crystals are used with  
modified waveform sampling  
electronics to reject pile-up  
events.



# Belle to Belle II

## Vertex detector

4 SVD layer  $\rightarrow$  2 layers  
DEPFET + 4 layers DSSD  
Expected resolution of  
 $\sim 25\mu\text{m}$  while in Belle  $\sim 50\mu\text{m}$

## CDC

Larger volume drift chamber,  
smaller drift cell. Faster  
electronics

## PID


More compact. Time of  
Propagation (barrel) and prox.  
foc. ARICH (Endcap) is used.  
Provide better  $K/\pi$  separation.

## KLM

Resistive place counter  
(Barrel)  
Scintillation + WLSF + MCCP  
(endcap)

## ECL

Old crystals are used with  
modified waveform sampling  
electronics to reject pile-up  
events.




ECL crystals and part of KLM sub-detector are re-used.



# Recent status









Currently we are running Phase 2, all sub-detectors are in except full vertex detector.

First collision on 26 April 2018







# Validation : Belle II is working



# Re-discovery of “November revolution”

in June



# Starting from the start: X(3872)

Most probable explanation:


Molecule with admixture of charmonium (*seems to be choice for now, others not ruled out yet*).

Precise Mass and Width studies.

- ✓ Expected yield of  $B^+ \rightarrow X(3872)(\rightarrow J/\psi\pi\pi)K^+ \sim 1500$  events (with  $10 \text{ ab}^{-1}$ )<sup>§</sup>
- ✓ Current yield of  $B^+ \rightarrow \psi'(\rightarrow J/\psi\pi\pi)K^+$  is  $\sim 3600$  events (at Belle).



Belle II should be able to observe

$$X(3872) \text{ or } \chi_{c1}' \rightarrow \chi_{c1}\pi^+\pi^-$$



Belle, PRD 81, 031103 (2010)

Informative to study  $X(3872) \rightarrow \overline{D}^0 D^{*0}$  in Belle II data



$$\text{Mass} \rightarrow 3872.9^{+0.6}_{-0.4} \text{ MeV}/c^2$$

<sup>§</sup>1/5 of total data

# Decays of X(3872)

## Measuring ratios of radiative decays

$$\mathcal{B}(X(3872) \rightarrow \psi' \gamma) / \mathcal{B}(X(3872) \rightarrow J/\psi \gamma) = 3.5 \pm 1.4 \quad \text{BaBar, PRL 102, 132001 (2009)}$$

$$< 2.1 \text{ (@90\% CL)} \quad \text{Belle, PRL 107, 091803 (2011)}$$

$$= 2.46 \pm 0.64 \pm 0.29 \quad \text{LHCb, NPB 886, 665 (2014)}$$

Expected yield of  $B^+ \rightarrow X(3872)(\rightarrow J/\psi \gamma) K^+ : \sim 400$  events (with  $10 \text{ ab}^{-1}$ )

Measure the above mention ratio precisely in order to constraint the admixture.

## Charged partner of X(3872)


Belle, PRD 84, 052004 (2011)

Negative search

$$\mathcal{B}(B^0 \rightarrow X(3872)^+ K^-) / \mathcal{B}(X(3872)^+ \rightarrow J/\psi \pi^+ \pi^0) < 4.2 \times 10^{-6}$$

If found, will be very promising for the tetraquark picture.

Absence of charged partner suggest X(3872) to be an iso-singlet state.




Suggests  $X(3872) \rightarrow J/\psi \pi^+ \pi^-$  is iso-spin violating decay ?

Belle and BaBar measured the allowed  $X(3872) \rightarrow J/\psi \pi^+ \pi^- \pi^0$

$$\frac{\mathcal{B}(X(3872) \rightarrow J/\psi \omega (\rightarrow \pi^+ \pi^- \pi^0))}{\mathcal{B}(X(3872) \rightarrow J/\psi \pi^+ \pi^-)} = 0.8 \pm 0.3$$


Belle II should measure this ratio.

BaBar, PRD 82 011101 (R) (2010)




# Production of X(3872)

Belle, PRD 97, 012005 (2018)



Measuring Absolute  $\mathcal{B}(B \rightarrow X(3872)K^+)$  will help in measuring  $\mathcal{B}(X(3872) \rightarrow \text{final state})$ .

Measurement is **“only possible at B factories”**  
(operating at center-of-mass energy of  $\Upsilon(4S)$  which decays into  $B\bar{B}$  pairs)



Missing mass recoiling against  $K^+$


$$M_{miss} = \sqrt{(p_{e^+e^-}^* - p_{tag}^* - p_h^*)^2}$$

$$\mathcal{B}(B^+ \rightarrow X(3872)K^+) < 2.6 \times 10^{-4} \text{ (@ 90\% CL)}$$

Belle II might measure this value.

➤ Not only for X(3872), but also for other states.

Belle, PRD91, 051101 (R) (2015)



$K^*(892)^0$  component in  $(K\pi)$  system in  $X(3872)$  does not dominate, *“in marked contrast”* to  $\psi'$  case.

With  $10 \text{ ab}^{-1}$ , Belle II will measure this precisely.

*Events will be similar to what we have now for  $\psi'$ .*



# Other production

Two photon processes

Study of  $\chi_{c2}(3930)$  using  $\gamma\gamma \rightarrow Z(3930) \rightarrow D\bar{D}$

Mass and width precision study.

$X(3915)$  (thought to be  $\chi_{c0}(2P)$ ) was

discovered in two photon process.

Currently,  $\chi_{c0}(2P)$  has been suggested to

be recently found  $X(3860)$  in  $J/\psi D\bar{D}$ .

Belle observed  $X(4350)$  in  $\gamma\gamma \rightarrow J/\psi\phi$ .


Recently, LHCb did amplitude analysis of

$B \rightarrow J/\psi\phi K$ , found several structures  $Y(4140)$ ,


$Y(4274)$ ,  $X(4500)$ ,  $X(4700)$  but not  $X(4350)$  (?)

Belle II should revisit with more data.

Belle, PRL 96 082003 (2006)



BaBar, PRD81 092003 (2010)



$X(3915)$  (thought to be  $\chi_{c0}(2P)$ ) was

discovered in two photon process.

Currently,  $\chi_{c0}(2P)$  has been suggested to

be recently found  $X(3860)$  in  $J/\psi D\bar{D}$ .



Belle observed  $X(4350)$  in  $\gamma\gamma \rightarrow J/\psi\phi$ .

Recently, LHCb did amplitude analysis of


$B \rightarrow J/\psi\phi K$ , found several structures  $Y(4140)$ ,

$Y(4274)$ ,  $X(4500)$ ,  $X(4700)$  but not  $X(4350)$  (?)


Belle II should revisit with more data.




Belle, PRL 104, 112004 (2010)




LHCb, PRD 95, 012002 (2017)



Belle, PRL 198, 082001 (2007)




Belle, PRL 100, 202001 (2008)



Double charmonium production, another interesting process through which Belle II can access  $C=+$  even states.

## $e^+e^- \rightarrow J/\psi\pi^+\pi^-$ study


# $\Upsilon$ ISRs



$$\frac{BR[Y(4260) \rightarrow Z(3895)^\pm \pi^\mp]}{BR[Y(4260) \rightarrow J/\psi \pi^+ \pi^-]} = (29.0 \pm 8.9)\%$$

- Belle II will compliment BESIII here.
- Expects improvement in mass resolution due to longer CDC
- One possible study  $e^+e^- \rightarrow \Upsilon(\rightarrow J/\psi \pi^0 \pi^0) \gamma|_{SR}$  for neutral partner

159±49±7 events





### Measured properties

- Mass = (3894.5±6.6± 4.5) MeV
- Width = (63±24±26) MeV

## $e^+e^- \rightarrow \psi' \pi^+ \pi^-$ study

Belle, PRD 91, 112007 (2015)




Mass = (4054±3± 1) MeV  
Width = (45±11±6) MeV

Any relation to  $Z(4050)^+ \rightarrow \chi_{c1} \pi^+$  ?  
Search  $Z(4430)^+ \rightarrow \psi' \pi^+$  as in  
 $B^0 \rightarrow \psi' \pi^+ K^-$  ?


Search for  $Z_{CS}^+$  in  $e^+e^- \rightarrow J/\psi KK$ .  
Study  $e^+e^- \rightarrow D^0 D^- \pi^+$  and  $e^+e^- \rightarrow \Lambda_c^+ \Lambda_c^-$ .

# Z : “with a charge”


Belle, PRD 88, 074026 (2013)



Belle, PRD80, 031104 (2009)



Belle, PRD80, 031104 (2009)



$(D^* D^*)^+$

Name	M (MeV/c <sup>2</sup> )	Γ (MeV)	J <sup>P</sup>	Process	
$Z_1(4050)^+$	$4051^{+24}_{-43}$	$82^{+51}_{-55}$	??	$B \rightarrow (\chi_{c1} \pi^+) K^-$	
$Z(4200)^+$	$4196^{+35}_{-37}$	$370^{+99}_{-149}$	1 <sup>+</sup>	$B \rightarrow (J/\psi \pi^+) K^-$	
$(D_0 D_1)^+$	$Z_2(4250)^+$	$4248^{+185}_{-45}$	$177^{+321}_{-72}$	??	$B \rightarrow (\chi_{c1} \pi^+) K^-$
$(D_1 D^*)^+$	$Z(4430)^+$	$4477 \pm 20$	$181 \pm 31$	1 <sup>+</sup>	$B \rightarrow (\psi' \pi^+) K^-$ $B \rightarrow (J/\psi \pi^+) K^-$

- Perform Dalitz analyses with more statistics: help in measuring and understanding these states with precision.
- At Belle II, search for new states using  $B^0 \rightarrow (\chi_{c2} \pi) K^+$  decay mode.
  - At  $10 \text{ ab}^{-1}$ , yield comparable to current Belle yield of  $B^0 \rightarrow (\chi_{c1} \pi) K^+$
- Possible study of  $B^0 \rightarrow (c \bar{c}) \pi^0 K^+$  in search for neutral partners.


# Bottomonium at Belle

Bottomonium spectrum is significantly different from charmonium spectrum.  
 $Z_b$  states were found in the  $\Upsilon(5S)$  decays and were clear signature of *eXotic* state.


Production ratio Belle, PRL 108 032001 (2012)

$$\frac{\Gamma(\Upsilon(5S) \rightarrow h_b(nP)\pi^+\pi^-)}{\Gamma(\Upsilon(5S) \rightarrow \Upsilon(2S)\pi^+\pi^-)} = \begin{cases} 0.45 \pm 0.08^{+0.07}_{-0.12} & \text{for } h_b(1P) \\ 0.77 \pm 0.08^{+0.22}_{-0.17} & \text{for } h_b(2P) \end{cases}$$



Decay to  $h_b$  should be suppressed due to spin flip!  
 $\Upsilon(5S) \rightarrow h_b(nP)\pi^+\pi^-$  decay mechanism seems to be *eXotic*



Fit  $MM(\pi)$  in  $M(h_b\pi)$  bins Belle, PRL 108, 122001 (2012)



Resonant structure of  $\Upsilon(5S) \rightarrow \Upsilon(nS)\pi^+\pi^-$




More precise measurements.

# More on $Z_b$

$B^+ \rightarrow J/\psi K^+$ ,  $B^+ \rightarrow D^0 (\rightarrow K^+ \pi) \pi^+$ ,  $B^+ \rightarrow D^0 (\rightarrow K^+ \pi \pi^+ \pi) \pi^+$ ,  
 $B^0 \rightarrow J/\psi K^0 (\rightarrow K^+ \pi)$ ,  $B^+ \rightarrow D^- (\rightarrow K^+ \pi^+ \pi) \pi^+$ ,  $B^0 \rightarrow D^{*-}$   
 $(\rightarrow D^0 [ \rightarrow K^+ \pi ] \pi) \pi^+$ ,  $B^0 \rightarrow D^{*-} (\rightarrow D^0 [ \rightarrow K^+ \pi \pi^+ \pi ] \pi) \pi^+$  and  
 $B^0 \rightarrow D^{*-} (\bar{D}^0 [ \rightarrow K^+ \pi \pi^0 ] \pi) \pi^+$


$\Upsilon(5S) \rightarrow B^* \bar{B}^{(*)} \pi$

Masses of  $Z_b(10610)^+$  and  $Z_b(10650)^+$  close to  $BB^*$  and  $B^* \bar{B}^*$  threshold



One B is fully reconstructed

$$rM(B\pi) = \sqrt{E_{cms}^2 - P_{B\pi}^2}$$





B is combined with  $\pi$  and recoil mass to  $(B\pi)$  combination is calculated

Belle, PRL116, 212001 (2016)



- $Z_b(10610)^+$  in  $BB^*$  and  $Z_b(10650)^+$  seen in  $B\bar{B}^*/B^*\bar{B}^*$ .
- $B^{(*)}B^*$  dominant mode of  $Z_b$  decays.

Belle II can confirm  $Z_b$  relation to  $B^{(*)}B^*$ .



Neutral  $Z_b^0$  in  $\Upsilon(5S) \rightarrow \Upsilon(nS)\pi^0\pi^0$

Belle, PRD88, 052016 (2013)




Belle II can study neutral  $Z_b^0$  and confirm in other modes also.




# Energy scan

- Many quarkonium-like states were found in energy scans in *ISR*,  $Y(4008)$  and  $Y(4260)$  in  $J/\psi\pi^+\pi^-$ ,  $Y(4360)$  and  $Y(4660)$  in  $\psi'\pi^+\pi^-$ ,  $\psi(4050)$  and  $\psi(4160)$  in  $J/\psi\eta$ .
  - Peaks observed in the cross-section depend on final state.
- Recent energy scan of  $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$  ( $n=1,2,3$ ) cross sections by Belle, show situation is different in bottomonium-like states.
  - All of cross-sections exhibits peaks at  $\Upsilon(10860)$  and  $\Upsilon(11020)$  resonances that are also seen in total hadronic cross sections.




## Energy scan of $e^+e^- \rightarrow h_b(nP)\pi^+\pi^-$ ( $n=1,2$ )

Belle, PRL 117, 142001 (2016)



Data consist of five energy points in  $\Upsilon(6S)$

- Evidence that proceed via intermediate  $Z_b$  state.
  - Only  $Z_b(10610)$  (excluded  $3.3\sigma$ )
  - Only  $Z_b(10650)$  produced not excluded significantly.



Current statistics is limited and Belle II will play crucial role here.

# Transition from $\Upsilon(5,6S)$ to molecular states

With unique data set at  $\Upsilon(6S)$ , Belle II can understand the  $\Upsilon(6S) \rightarrow Z_b$  decay


$$\Upsilon(6S) \rightarrow h_b(nP) \pi^+ \pi^-, \Upsilon(mS) \pi^+ \pi^- [n=1,2 ; m=1,2,3]$$

If  $Z_b$  molecular state, then Heavy Quark Spin symmetry suggest there should be 2/4 molecular partner bottomonium-like state ( $W_b$ )

$$\Upsilon(5S,6S) \rightarrow W_{b0} \gamma$$

$$\Upsilon(6S) \rightarrow W_{b0} \pi^+ \pi^-$$

$$W_{b0} \rightarrow \eta_b \pi, \chi_b \pi, \Upsilon \rho$$



Voloshin, PRD 84, 031502(R)(2011)

# Future summary

- Quarkonium sector is not as simple as one expects.
- Many new states have been found with puzzling nature.
- Still not fully understood in spite of the best efforts by all the experiments.
- Belle II will play an important role along with LHCb and BESIII to understand them.
- Belle II detector already started collecting data and hope to provide fruitful results soon.





**Thank you**