


Search for Dark Particles at Belle and Belle II

Igal Jaegle

University of Florida

for the Belle/Belle II Collaborations

Dark Interactions, October 4 - 7, 2016
Brookhaven National Laboratory



UF UNIVERSITY of FLORIDA



Table of contents

- 1 Introduction
- 2 Setup
- 3 Search for a dark vector gauge boson
- 4 Search for the dark photon and the dark Higgs boson
- 5 Search for the dark photon and light dark matter
- 6 Conclusion

Search for new hidden forces / Introduction

Dark matter represents about 80% of all matters

- Dark matter naturally explained by Supersymmetry, but
- Absence of Supersymmetry in LHC gives new light to Dark Sector Models
- Search for new hidden forces accessible to Belle

PRD 75 115017 (2007)



PRD 89 114008 (2014)

- ▶ Coupling of charged matter to new, dark photon, A' , $q = \epsilon e$

▶ $\mathcal{L} = -\frac{1}{2}\epsilon F_{dark}^{\mu\nu} F_{\mu\nu}$

- ▶ Coupling of all quarks to new, baryonic boson, U' , $g_{U'} = \sqrt{4\pi\alpha_{U'}}$

▶ $\mathcal{L} = \frac{1}{3}g_{U'}\bar{q}\gamma^\mu q U'_\mu$



- ▶ Feynman diagram
- ▶ A' short or long lived or invisible



- ▶ Feynman diagrams for hadronic U' decay
- ▶ Above π threshold U' short lived

All models point to new particles with mass of the order of MeV – GeV

Search for new hidden forces / Introduction

Dark matter represents about 80% of all matters

- Dark matter naturally explained by Supersymmetry, but
 - Absence of Supersymmetry in LHC gives new light to Dark Sector Models
 - Search for new hidden forces accessible to Belle
- Branching ratio for A' decay [PRD 79 115008 \(2008\)](#)
- Branching ratio for U' decay [PRD 89 114008 \(2014\)](#)




All models point to new particles with mass of the order of MeV – GeV

KEKB and SuperKEKB

KEKB/SuperKEKB collider, located in Japan, Tsukuba, is the world's highest-luminosity electron-positron collider


- 1999-2010: Belle collected $\mathcal{L}_{int} = 1050 \text{ fb}^{-1}$ at $\Upsilon(1S, 2S, 3S, 4S, 5S)$ and continuum
- 2016-2026: Belle II (upgrade version of Belle) expects to collect $\mathcal{L}_{int} = 50 \text{ ab}^{-1}$



Schematic view of KEKB

$$L = \frac{\gamma_{\pm}}{2e r_e} \left(1 + \frac{\sigma_y^*}{\sigma_x^*} \right) I_{\pm} \xi_{y\pm} \left(\frac{R_L}{R_{\xi}} \right)$$

Lorentz factor γ_{\pm}
 Beam current I_{\pm}
 Beam-Beam parameter $\xi_{y\pm}$
 Geometrical reduction factors (crossing angle, hourglass effect) $\left(\frac{R_L}{R_{\xi}} \right)$
 Beam aspect ratio at IP $\left(1 + \frac{\sigma_y^*}{\sigma_x^*} \right)$
 Vertical beta function at IP $\beta_{y\pm}^*$
 Minimum value is limited by hourglass effect



S. Kurokawa and E. Kikutani, NIM A 499, 1 (2003)

KEKB beam vs. SuperKEKB nano-beam

$$\text{Belle II } \mathcal{L}_{peak} = 8 \times 10^{35} \text{ cm}^{-2} \text{ s}^{-1}$$


Belle and Belle II experiments

CP violation measurement in the B-meson system with Belle and *BABAR*, established the Kobayashi Maskawa mechanism as a valid description of CP violation in the Standard Model.

● Main motivations

- ▶ Study of CP violation (i.e. matter-antimatter asymmetry)
- ▶ Study of heavy flavor
- ▶ Search for physics beyond the Standard Model

● Complementary to efforts at energy frontier





Belle II is an upgrade of Belle

Search for a dark vector gauge boson

New Belle results submitted to PRD: [arxiv:16090.5599](https://arxiv.org/abs/16090.5599), analysis and article of Eunil Won

- Search for $U' \rightarrow \pi^+\pi^-$ in $D^0 \rightarrow K_S^0\eta$, $\eta \rightarrow U'\gamma$ using $977fb^{-1}$ of Belle data
- Exclusive charm meson decays to reduce background





▶ $K_S^0\eta$ invariant mass

▶ D^*-D^0 mass difference

- Cut on D^0 and mass difference and look at $\pi^+\pi^-\gamma$ invariant mass

Search for a light vector gauge boson


- Background estimated and subtracted using the side bands



- ▶ $\pi^+\pi^-\gamma$ invariant mass
- ▶ $\pi^+\pi^-$ invariant mass after background subtraction
- $\pi^+\pi^-$ -line shape fitted by $\frac{d\Gamma}{ds} \propto |P(s)F_V(s)|^2(m_\eta^2 - s)^3s(1 - 4m_\pi^2/s)^{3/2}$
[PLB 707, 243 (2012) and PLB 707, 184 (2012)]
 - ▶ $P(s)$ reaction-specific perturbative part
 - ▶ $F_V(s)$ pion vector form factor
- **No signal found**, example of U' MC sim. signal of 400 MeV/c² mass is shown

New Belle limit

Submitted to PRD, arxiv:16090.5599 (Eunil Won)




- 95% CL limit on the baryonic fine structure constant
- Better limit for $m_{U'} > 450 \text{MeV}/c^2$ and $\phi \rightarrow e^+ e^- \gamma$

Search for the dark photon and the dark Higgs boson

Belle limits, PRL 114 211801 (2015). Production in the so-called Higgs-strahlung channels, $e^+e^- \rightarrow A'h'$, with $h' \rightarrow A'A'$.

- A' and h' assuming prompt decays
- $m_{h'} > 2m_{A'}$
- $0.1 < m_{A'} < 3.5 \text{ GeV}/c^2$ and $0.2 < m_{h'} < 10.5 \text{ GeV}/c^2$



α_D : dark sector constant

ϵ : kinetic mixing

- 10 exclusive channels: $3(l^+l^-)$, $2(l^+l^-)(\pi^+\pi^-)$, $2(\pi^+\pi^-)(l^+l^-)$, and $3(\pi^+\pi^-)$, where l^+l^- is an electron or muon pair
- 3 inclusive channels for $m_{A'} > 1.1 \text{ GeV}/c^2$: $2(l^+l^-)X$, where X is a dark photon candidate detected via missing mass


If $\alpha_D = 1$, Higgs-strahlung channels most sensitive to A'

Belle limits / results

● Belle limits for $\mathcal{L} = 977 \text{ fb}^{-1}$ on $\mathcal{B} \times \sigma_{\text{Born}}$ and σ_{Born}

▶ 90% CL upper limit for each of the 13 final states

▶ 90% CL upper limit on the combined Born cross section




● 90 % Credibility Level (CL) upper limit determined by Bayesian inference method with the use of Markov Chain Monte Carlo *A'. Caldwell et al., CPC 180 (2009) 2197-2209*

Limits from $3(\pi^+\pi^-)$ and $2(e^+e^-)X$ are the first placed by any experiment

Limits on the product of $\alpha_D \epsilon^2$ / results

Belle combined limits compared to *BABAR* combined limits

- Belle limits for $\mathcal{L} = 977 \text{ fb}^{-1}$ based on the Born cross section, ISR effect non negligible
- *BABAR* limits for $\mathcal{L} = 520 \text{ fb}^{-1}$ based on the visible cross section [PRL 108 211801 \(2012\)](#)



90% CL upper limit on the product $\alpha_D \times \epsilon^2$ versus dark photon mass (top row) and dark Higgs boson mass (bottom row)

- Assuming branching fractions and couplings versus cross section from [B. Batell et al. PRD 79 \(2009\) 115008](#)


Results scale nearly linearly with integrated luminosity. This bodes well for future searches with Belle II.

Belle II prospects for the Higgs-strahlung channels

Predicted Belle II upper limits $U_{\alpha_D \varepsilon^2}$ in the $\alpha_D \varepsilon^2$ vs $m_{A'}$ vs $m_{h'}$ plane by scaling the Belle limits linearly with the integrated luminosity:

$$\frac{U_{\alpha_D \varepsilon^2}}{U_{\alpha_D \varepsilon^2}^0} = \frac{\mathcal{L}^0}{\mathcal{L}},$$



where the superscript 0 corresponds to Belle values. \mathcal{L} is integrated luminosity. The scaling uses both statistical and systematic uncertainties.



Belle (II) prospect for the radiative decays

Predicted Belle II upper limits extrapolated from *BABAR* [PRL 113, 201801 \(2014\)](#) (C. Hearty, B2TIP2014)

- $e^+e^- \rightarrow \gamma A'$, $A' \rightarrow l^+l^-$, with $l = e$ or μ
- Belle II will have an improve low multiplicity trigger compared to Belle





Left: Belle II prediction. Right: preliminary Belle and Belle II preliminary fiducial detection efficiency for $A' \rightarrow \mu^+\mu^-$

Belle II will have a better efficiency for low momentum muon compared to Belle

Belle (II) prospect for the radiative decays

Predicted Belle II upper limits extrapolated from *BABAR* [PRL 113, 201801 \(2014\)](#) (C. Hearty, B2TIP2014)

- $e^+e^- \rightarrow \gamma A'$, $A' \rightarrow I^+I^-$, with $I = e$ or μ
- Belle II dimuon invariant mass resolution improved by $\sim 35\%$ compared to Belle





Left: Belle II prediction. Right: simulation for $m_{A'} = 5.015 \text{ GeV}/c^2$

Complementary to fixed target experiments

Belle (II) prospect for the radiative decays

Predicted Belle II upper limits extrapolated from *BABAR* [arxiv:0808.0017](https://arxiv.org/abs/0808.0017) (C. Hearty, B2TIP2014)

- $e^+e^- \rightarrow \gamma A'$, $A' \rightarrow \chi\chi$, χ light dark matter R. Essig et al. [arXiv:1309.5084](https://arxiv.org/abs/1309.5084)
- Require implementation of a single photon trigger in Belle II



Left: Belle II prediction. Right: Simulated mono-energetic photon signature for $m_{A'} = 6 \text{ GeV}/c^2$

Belle II is expected to have a better single photon trigger with a lower energy sum than *BABAR*

Conclusion

- New Belle limit has been presented on the baryonic fine constant, $\alpha_{U'}$, in a exclusive charm decays
 - ▶ $280 < m_{U'} < 550 \text{ MeV}/c^2$
 - ▶ We found that:
 - ★ No signal found
 - ★ Better limit for $m_{U'} > 450 \text{ MeV}/c^2$ and $\phi \rightarrow e^+e^-\gamma$
- Belle limits for prompt decays of the dark photon and the dark Higgs boson:
 - ▶ $0.1 < m_{A'} < 3.5 \text{ GeV}/c^2$
 - ▶ $0.2 < m_{h'} < 10.5 \text{ GeV}/c^2$
 - ▶ We found that:
 - ★ No significant excess over the background estimation
 - ★ Belle limit improvement scales nearly linearly with integrated luminosity
- Ongoing Belle analysis on $e^+e^- \rightarrow A'\gamma$ (prompt and displaced vertex), $e^+e^- \rightarrow \chi\chi\gamma$, $e^+e^- \rightarrow \mu^+\mu^-Z'$, and $e^+e^- \rightarrow \tau^+\tau^-h''$
- Belle II will also search for dark particles
- With 50 ab^{-1} , Belle II might potentially also cross-check any signals discovered by fixed target experiments
- First collisions in 2017 with partial vertex detector

Thank you for your attention