

Search for a dark photon and a visible long-lived dark higgs boson at the Belle II experiment

La Thuile 2025 - Les Rencontres de Physique de la Vallée d'Aoste, 9-15 March 2025

Foteini Faidra Trantou^{1,2}, on behalf of the Belle II collaboration

¹University of Pisa

²INFN sezione di Pisa

foteini.trantou@pi.infn.it



Outline

- Introduction
 - The Dark Sector
 - Analysis overview
- Preliminary studies
 - Signal generation
 - Background suppression
 - Results
- Summary

Introduction

The Dark Sector at Belle II

■ Intensity frontier experiment

- Operating at the SuperKEKB accelerator in Japan, an e^+e^- asymmetric energy collider
- Current data taking status: Recorded a total integrated luminosity of more than **575 fb⁻¹**.



➤ Provides a unique opportunity to search for new physics such as dark matter (DM)

DM could be part of a larger hidden/dark sector, interacting with Standard Model (SM) matter **very weakly via subtle mixing through *portals***.

➤ *High sensitivity to dark sectors*: Can access the mass range naturally favored by light dark sectors at $M_{\text{Dark matter}} \sim \mathcal{O}(\text{MeV} - \text{GeV})$

➤ *Large potential for dark matter discoveries*

High luminosity, dedicated triggers and excellent reconstruction capabilities for low multiplicity final states and missing energy signatures



See talks by [M.Veronesi](#) and [M.Campajola](#) for more information on Belle II searches


The Dark Higgsstrahlung process

- Exploring two dark sector particles: the **dark photon A'** and the **dark Higgs boson h'**

The mass of the A' can arise, in close analogy with the SM, via a spontaneous symmetry breaking mechanism

→ Introduces a new scalar particle: a dark Higgs boson h'

- The A' interacts with the SM photon through a kinetic mixing mechanism with strength ϵ while h' couples to A' with α_D
- Different scenarios depending on the mass hierarchy
 - $m_{h'} > m_{A'}$: $h' \rightarrow A'A'$ with up to 6 tracks in the final state
Investigated by [BaBar](#) (2012) and [Belle](#) (2015)
 - $m_{h'} < m_{A'}$: h' decays outside the detector resulting invisible
Constrained by [KLOE](#) (2015) and [Belle II](#) (2023)



Analysis overview

- Focusing on the light dark higgs h' scenario ($m_{h'} < m_{A'}$)

Considered model: h' is naturally the lightest dark sector state and decays into SM particles via interacting with the SM Higgs at a **mixing angle θ**


- Search for a dark photon A' and a visible long lived dark Higgs boson h' via the *dark Higgsstrahlung process*:

$$e^+e^- \rightarrow A'(\rightarrow \mu^+\mu^-)h'(\rightarrow \mu^+\mu^-/h^+h^-) \text{ where } h = \pi, K$$

- **Motivation to search for a long-lived dark higgs boson**

- θ affects the lifetime of the dark Higgs: $\tau_{h'} \propto 1/\sin^2(\theta)$
Renders it naturally a **long-lived particle** for small values of θ

→ **Search:** h' decaying **inside** the Belle II detector



Key model parameters

- Mass of the **dark photon** $m_{A'}$
- Mass of the **dark higgs** $m_{h'}$
- Kinetic mixing with **strength ϵ** of the A' with SM
- **Mixing angle θ** of the h' with the SM higgs
- **α_D** the coupling between the h' and the A'

[Duerr M. et al. J. High Energy. Phys. 2021, 146 (2021)]

Analysis overview

- Focusing on the light dark higgs h' scenario ($m_{h'} < m_{A'}$)

Considered model: h' is naturally the lightest dark sector state and decays into SM particles via interacting with the SM Higgs at a **mixing angle θ**


- Search for a dark photon A' and a visible long lived dark Higgs boson h' via the *dark Higgsstrahlung process*:

$$e^+e^- \rightarrow A'(\rightarrow \mu^+\mu^-)h'(\rightarrow \mu^+\mu^-/h^+h^-) \text{ where } h = \pi, K$$


- **Motivation to search for a long-lived dark higgs boson**

- θ affects the lifetime of the dark Higgs: $\tau_{h'} \propto 1/\sin^2(\theta)$
Renders it naturally a **long-lived particle** for small values of θ

→ **Search:** h' decaying **inside** the Belle II detector



Existing limits to my model from other dark scalar searches



L. Adachi et al. (Belle II Collaboration),
Phys. Rev. D 108 (2023)

Preliminary studies

Signal signature and reconstruction


- Simultaneous search of a dark photon A' and a dark higgs h' in **four-track** events and final states with **no missing energy** E_{miss}
- **2 prompt tracks**, associated with the A' decaying into a pair of **muons** + **2 tracks forming a displaced vertex (LLP h' decay)**

Signal signature is two narrow peaks, at different invariant masses.

- **Current focus on $h' \rightarrow \mu^+ \mu^-$** :
- Signal generation using MadGraph5 aMC@NLO with Initial State Radiation (ISR) effects included.
- The h' production vertex is displaced according to the lifetime in **ct** [cm].

Event
pre-selections

- **Maximum 6 tracks** in the final state coming from the interaction point (IP), with **two tracks** identified as **muons**
- **No missing energy** with $m_{\text{Recoil}} < 6.5 \text{ GeV}/c^2$



Signal efficiency and resolutions

Reconstruction efficiency as a function of the dark Higgs lifetime

Sensitivity to long-lived particles with notable ϵ_{reco} even at extended lifetimes



■ To estimate the resolutions, we fit the signal shape with a Double Sided Crystal Ball function with the following characteristics:

- Gaussian core and exponential tails on each side
- Fixed mean according to the respective invariant mass.

■ We extract the width of the signal peaks and obtain the resolutions:

→ σ increases as a function of the invariant masses but remains small at $\sigma_{A'} = 4 - 40 \text{ MeV}/c^2$ and $\sigma_{h'} = 1 - 14 \text{ MeV}/c^2$.

■ Defining the signal mass windows : Window size is set $\pm 5 \sigma$ around the nominal value of the corresponding h' and A' mass.



Background studies


Characterization and suppression

Main SM background contributions from combinatorial $e^+e^- \rightarrow q\bar{q}$ events, **two** and **four** lepton events such as $e^+e^- \rightarrow \mu^+\mu^-(\gamma)$.



- *Applying preselection* leads to **96.1 %** background suppression (no mass selections here)

- Using run dependent samples, each rescaled to the target dataset size of this analysis:

$$\int \mathcal{L} dt = 361 fb^{-1}$$



Accessible search region has a **triangle shape** corresponding to the mass plane $m_{h'} - m_{A'}$ in signal events.







Background suppression

Utilizing a collection of **discriminating variables**, motivated by signal kinematics:

- **dr** : Transverse distance of the h' decay vertex with respect to the IP [cm]
- **missing_** E_{CMS} : Total missing energy of the four reconstructed tracks
- **$\theta_{\text{point}}^{h'}$** : Pointing angle of the dark higgs h'
- **θ_{helicity}** : Helicity angle of the dark photon A' and the dark higgs h'

Here: Mass selections are applied, both for signal and background, at $\pm 5 m_{\text{inv}}$







Background suppression

Utilizing a collection of **discriminating variables**, motivated by signal kinematics:

- **dr** : Transverse distance of the h' decay vertex with respect to the IP [cm]

Sufficient displacement of h' - LLP analysis :
Common cut on across the mass plane at $0.2 < dr < 110 \text{ cm}$





Background accumulation

Studied via the correlation of the discriminant variables



- $\text{missing_E}_{\text{CMS}}$ vs $\theta_{\text{point}}^{h'}$:

Background peaks for **high values** of the missing energy and for **non pointing** h' vertices



- $\theta_{\text{helicity}}^{A'}$ vs $\theta_{\text{helicity}}^{h'}$:

Background distribution peaks at ± 1 as muons do not originate from a common source



Background accumulation





Across the mass plane, we observe:

- ❖ Different background contributions per region

$\tau^+\tau^-$ and $q\bar{q}$ events populate mainly R2

- ❖ Uneven background distributions

Mostly in R3 where $N_{\text{bkg}}(\mu^+\mu^-)$ is 79.6% of the total events.






Event selection strategy

- Utilizing the **signal distribution shape**
- We apply a **box shaped cut** and perform a 2D **missing_** E_{CMS} **vs** $\theta_{\text{point}}^{h'}$ scan , followed by a scan on $\theta_{\text{helicity}}^{h'}$

➤ Selections are optimized in each region separately using the **Punzi**_{FOM}: $\frac{\epsilon_{sig}(t)}{\frac{a}{2} + \sqrt{N_{bkg}}}$

➤ **Strong suppression** achieved using particle identification also on the h' **muons**

- Negligible effect on the signal efficiency ϵ_{sig}
- Significant reduction for background



Signal efficiency and background

Combining all discriminant variables

Signal efficiency remains high across the whole mass plane with

$\epsilon_{\text{sig}} = 32.5\%$ and, after applying $\mu^{h'}_{\text{ID}}$, $\epsilon_{\text{sig}} = 28.5\%$



- $\epsilon_{\text{sig}} \sim 3\%$ even at extended lifetimes such as $c\tau_{h'} = 100$ cm

Expecting minimum background levels:

Selections result in high background rejection power

Remaining background, inside each signal window, can be very low and varies across the mass plane

Plan for signal yield extraction:
Cut and count technique



Summary

- Search for a dark photon A' and a visible long lived dark Higgs h' boson at the Belle II experiment via the *dark Higgsstrahlung process*:

$$e^+e^- \rightarrow A'(\rightarrow \mu^+\mu^-)h'(\rightarrow \mu^+\mu^-/h^+h^-) \text{ where } h = \pi, K$$

- *Dark higgs could interact with the SM higgs at a mixing angle θ*

For a **LLP h' decay scenario**, we can search h' decaying inside the Belle II detector

- Preliminary studies on $h' \rightarrow \mu^+\mu^-$ show:
 - Significant background suppression utilizing various discriminating analysis variables
 - Signal efficiency > **28%** across the mass plane
- **Ongoing:** Estimate the Belle II's sensitivity to the signal cross section for $h' \rightarrow \mu^+\mu^-$
- **Next steps:**
 - Include $h = \pi, K$ final states
 - Trigger and control data studies
 - Calculate and assign the systematic uncertainties

Thank you for your attention

Backup


The Belle II experiment

Overview

- Operating at the SuperKEKB accelerator (Tsukuba, JP), an e^+e^- asymmetric energy collider
- Optimized for the production of B meson pairs, but also D mesons and τ leptons.
- Collisions occur mainly at $\sqrt{s} = 10.58 \text{ GeV}$, corresponding to the $m_{\text{inv}} = Y(4S)$

Data taking status

- Run I (2019-2022) + Run II (February 2024-Now):
Recorded a total luminosity of more than **575 fb^{-1}** .
- SuperKEKB reached world's highest instantaneous luminosity at $\mathcal{L} = 5.1 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$.
- Target: $\int \mathcal{L} dt = 50 \text{ ab}^{-1}$ (**$50 \times$ Belle dataset!**)



Phases of the Experiment

- Belle II's data taking timeline is separated into three phases.
- ❖ **Phase (2016) + Phase II (April - July 2018) :**
Dedicated to commissioning both the machine and the detector.


Collected **0.5 fb⁻¹** of data with an incomplete vertex detector (VXD).

- ❖ **Phase III (March 25, 2019 - Now) :** Beginning of a full-scale data collection

- **Run I (2019-2022):** Recorded a total of **427 fb⁻¹**
- **Run II (January 2024-Now)**

The total luminosity collected by the experiment for


Run I and Run II amounts to **575.47 fb⁻¹**



Signal studies

Branching fractions

- Search can be performed in various final states, depending on the **decay of the dark Higgs h'**
- **Limited in mass for hadrons**
For pions and Kaons, current theoretical calculations provide predictions up to $m_{h'} = 2 \text{ GeV}/c^2$.
- **Other decays of the dark Higgs (not considered)**
 - Electron decay**
Existing limits below the **di-muon threshold** are quite strong, while above it, the e^- decay is heavily suppressed.
 - Tau decay (not depicted)**
Introduces complications in the reconstruction of the final state due to its neutrino decay and the no missing energy requirement we impose.




BR of the dark Higgs decaying into four different final states as a function of its mass.

Signal studies

Expected cross section

- Cross section of the process $e^+e^- \rightarrow A'(\rightarrow \mu^+\mu^-) h'(\rightarrow \mu^+\mu^-)$ evaluated by MadGraph at generator level
- **Dependency on the boson masses:**
Accessible search region has **triangle shape** that corresponds to the mass plane $m_{h'} - m_{A'}$ in signal events.
 - $\sigma \sim 10^{-2} - 10^{-1} \text{ fb}$ for $m_{h'} > 4 \text{ GeV}/c^2$
 - $\sigma \sim 10^2 \text{ fb}$ for $m_{h'} < 0.5 \text{ GeV}/c^2$
- **Dependency on other parameters**
The mixing angle and lifetime of the dark higgs are **related** via

$$\tau_{h'} \propto 1 / \sin^2(\theta)$$
 - σ is **stable** up to $\theta = 0.1 \text{ rad}$, with noticeable effects appearing at higher values.
 - σ shows **negligible** variation in relation to the h' lifetime.



For comparison: $\sigma(B\bar{B}) = 1.1 \text{ nb}$

Signal studies


Reconstruction efficiency

- Efficiency = #Correctly reconstructed events / #All generated events

as a function of the true transverse distance of the reconstructed h' decay vertex dr with respect to the IP

$$\epsilon_{\text{reco}} = \frac{h'_{\text{reconstructed}}}{h'_{\text{generated}}}$$

- Efficiency decreases as the distance of the h' decay vertex increases
- ϵ_{reco} greater for prompt h' decays, maintaining values above 40% for decay distances up to $dr = 25$ cm.




Using Monte Carlo truth matching for selecting signal events

Discriminating variables (1)

■ Transverse distance of the h' decay vertex dr

- Region close to the IP is expected to be dominated by SM backgrounds

→ Requiring that the h' decays with sufficient displacement at $0.2 < dr < 110 \text{ cm}$ (upper limit is set by the location of the CDC tracking detector at Belle II)




■ Total missing energy of the four reconstructed tracks

- The **signal** should peak at values close to zero

Deviations could derive from the ISR, misreconstruction or resolution effects.


- Background peaks at larger values.



Discriminating variables (2)


■ Pointing angle of the dark higgs h'

The angle between the **vector** connecting the reconstructed displaced vertex (DV) to the IP and the momentum vector \mathbf{p}_{DV} of the dimuon system for h' decay in signal events.



Depicted in $-\log_{10}(1 - \cos(\theta_{point}^{h'}))$:


- Expecting high values for **signal** since h' vertices are pointing back to the IP
- **Background** candidates are made up from random track combinations and peak at **low values**



Discriminating variables (3)

■ Helicity angle of the dark photon A' and the dark higgs h'

The angle between the flight direction of the h' in the CMS system and the μ^+ in the h' rest frame (same for the A')



- Signal is a resonance decaying into two fermion particles
→ Observing an **almost flat** distribution of $\cos(\theta_{\text{hel}})$ for A' and h'
- No direct physical meaning for the background:
muons do not originate from a common resonance
and the distribution peaks at ± 1

