



Quarkonium at Belle II


Speaker: Junhao Yin, on behalf of Belle II Collaboration



Junhao Yin, July 2021

outline

- Introduction
 - Quarkonium and the exotics
 - Questions
- The Belle II experiment
- Searching for exotics on Belle II
 - Re-discoveries with “early Phase III data”
 - Prospect
- Summary




In 1963, Murray Gell-mann proposed the quark model.



Hadrons with more than 3 quarks were predicted.

Junhao Yin, July 2021

Conventional hadrons




Non-standard hadrons



A simpler and more elegant scheme can be constructed if we allow non-integral values for the charges. We can dispense entirely with the basic baryon b if we assign to the triplet t the following properties: spin $\frac{1}{2}$, $z = -\frac{1}{3}$, and baryon number $\frac{1}{3}$. We then refer to the members $u^{\frac{2}{3}}$, $d^{-\frac{1}{3}}$, and $s^{-\frac{1}{3}}$ of the triplet as "quarks" q and the members of the anti-triplet as anti-quarks \bar{q} . **Baryons** can now be constructed from quarks by using the combinations **(qqq), (qqqq \bar{q}), etc.**, while **mesons** are made out of **(q \bar{q}), (qq \bar{q} \bar{q}), etc.** It is assuming that the lowest baryon configuration (qqq) gives just the representations **1, 8, and 10** that have been observed, while the lowest meson configuration (q \bar{q}) similarly gives just **1 and 8**.


Published in Physics Letters 8, 214 (1964);
 Similar idea by G. Zweig, CERN-TH-401 (1964).
QCD does not forbid hadrons with $N_{quark} \neq 2,3!!$

No solid evidence until 2003, the observation of $X(3872)$
 Since then, we have a golden era on the discovery of the exotic state.




Junhao Yin, July 2021


What are they?



What are they?



Belle II Detector




2nd Generation B-Factory Detector

- High-luminosity performance
- Much improved vertexing
- Novel Cherenkov PID (TOP)
- Other upgrades...

Belle II TDR, arXiv:1011.0352

Junhao Yin, July 2021



Corona-pandemic affected activity at KEKB

BUT

we are very active and data taking is ongoing!

Increasing by $1 \sim 1.5 \text{ fb}^{-1}$ per day

Luminosity record: $3.12 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$

Goal: $65 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$

Will accumulate around 50 ab^{-1} till 2026

Junhao Yin, July 2021

X(3872)

Most concerned particle in charmonium sector!

- Highest citation in Belle!
- Still many puzzles.
- decays ratios?
- lineshape?
- Real nature?

Observation of a narrow charmonium - like state in exclusive $B^{+-} \rightarrow K^{+-} \pi^+ \pi^- J/\psi$ decays #1

Belle Collaboration • [S.K. Choi](#) (Gyeongsang Natl. U.) et al. (Sep, 2003)



Published in: *Phys.Rev.Lett.* 91 (2003) 262001 • e-Print: [hep-ex/0309032](#) [hep-ex]

pdf links DOI cite

1,953 citations


Parameter index	Decay mode	Branching fraction
1	$X(3872) \rightarrow \pi^+ \pi^- J/\psi$	$(4.1^{+1.9}_{-1.1})\%$
2	$X(3872) \rightarrow D^{*0} \bar{D}^0 + c.c.$	$(52.4^{+25.3}_{-14.3})\%$
3	$X(3872) \rightarrow \gamma J/\psi$	$(1.1^{+0.6}_{-0.3})\%$
4	$X(3872) \rightarrow \gamma \psi(3686)$	$(2.4^{+1.3}_{-0.8})\%$
5	$X(3872) \rightarrow \pi^0 \chi_{c1}$	$(3.6^{+2.2}_{-1.6})\%$
6	$X(3872) \rightarrow \omega J/\psi$	$(4.4^{+2.3}_{-1.3})\%$
7	$B^+ \rightarrow X(3872) K^+$	$(1.9 \pm 0.6) \times 10^{-4}$
8	$B^0 \rightarrow X(3872) K^0$	$(1.1^{+0.5}_{-0.4}) \times 10^{-4}$
	$X(3872) \rightarrow \text{unknown}$	$(31.9^{+18.1}_{-31.5})\%$

Phys.Rev.D 100 (2019) 9, 094003



Phys.Rev.D 102 (2020) 9, 092005

Rediscovery of $X(3872)$




Reconstruction of final states

- $B^\pm \rightarrow \pi^+ \pi^- J/\psi (l^+ l^-) K^\pm$
- $B^0 \rightarrow \pi^+ \pi^- J/\psi (l^+ l^-) K_S$

Selection criteria





- Particle identification
- continuum suppression
- kinematics criteria: $M_{bc}, |\Delta E|$



$$M(\pi^+ \pi^-) > M(\pi^+ \pi^- l^+ l^-) - m(J/\psi) - 0.150 \text{ GeV}/c^2$$

Junhao Yin, July 2021

Rediscovery of $X(3872)$



Significant $\psi(2S)$
A good control sample

$$\frac{B^0 \rightarrow X(3872)K^0}{B^\pm \rightarrow X(3872)K^\pm} = 0.5 \text{ is assumed}$$

First $X(3872)$ at Belle II:

$$N_{\text{sig}} = 14.4 \pm 4.6$$

Evidence of $X(3872)$ with 4.6σ

Junhao Yin, July 2021

ISR preliminary studies

Goal:

- Exotic states with $J^{PC} = 1^{--}$ and their decays

Golden Channels	$E_{c.m.}$ (GeV)	Statistical error (%)	Related XYZ states
$\pi^+\pi^- J/\psi$	4.23	7.5 (3.0)	$Y(4008), Y(4260), Z_c(3900)$
$\pi^+\pi^- \psi(2S)$	4.36	12 (5.0)	$Y(4260), Y(4360), Y(4660), Z_c(4050)$
$K^+K^- J/\psi$	4.53	15 (6.5)	Z_{cs}
$\pi^+\pi^- h_c$	4.23	15 (6.5)	$Y(4220), Y(4390), Z_c(4020), Z_c(4025)$
$\omega\chi_{c0}$	4.23	35 (15)	$Y(4220)$

10 ab^{-1} 50 ab^{-1}

Competition:

- BESIII: $> 0.5 \text{ fb}^{-1}$ each point $\in [4.13, 4.70] \text{ GeV}$

Advantage:



- ISR is a “free energy scan”!
- Reach a really low energy region

Need really huge dataset!!

Junhao Yin, July 2021

ISR preliminary studies

- $e^+e^- \gamma_{\text{ISR}} \rightarrow \pi^+\pi^- J/\psi(\ell^+\ell^-)$ final states
 - Nominal PID requirements
 - $|M(J/\psi) - M(\text{PDG})| < 75 \text{ MeV}$
 - ISR photon not required (high efficiency)
 - $|MM^2(\pi^+\pi^- J/\psi)| < 2 \text{ GeV}^2$
- Clear observation of ISR $\psi(2S)$ signals
- Next step: “Y(4260)” rediscovery
 - Expect ~ 60 total events per 100 fb^{-1}




Junhao Yin, July 2021

Bottomonia at Belle II

$\Upsilon(5,6S)$:

- Study of Z_b : branching ratios, decays...
- Search for new/predicted resonances
- $\Upsilon(5,6S)$ transitions are different
 - hint for non $b\bar{b}$ nature?

An extra resonance around 10.750 GeV?




Junhao Yin, July 2021

Bottomonia at Belle II

$\Upsilon(5,6S)$:

- Study of Z_b : branching ratios, decays...
- Search for new/predicted resonances
- $\Upsilon(5,6S)$ transitions are different
 - hint for non $b\bar{b}$ nature?

An extra resonance around 10.750 GeV?




Junhao Yin, July 2021

Bottomonia at Belle II

$\Upsilon(5,6S)$:

- Study of Z_b : branching ratios, decays...
- Search for new/predicted resonances
- $\Upsilon(5,6S)$ transitions are different
 - hint for non $b\bar{b}$ nature?


An extra resonance around 10.750 GeV?



Junhao Yin, July 2021

JHEP 10 (2019) 220

Dipion transition among bottomonia




Study $e^+e^- \rightarrow \pi^+\pi^-\mu^+\mu^-$ (+ γ undetected)

- $Y(4S) \rightarrow \pi^+\pi^- Y(nS)$
- $e^+e^- \rightarrow \gamma_{\text{ISR}} Y(mS), Y(mS) \rightarrow \pi^+\pi^- Y(nS)$

Junhao Yin, July 2021

Dipion transition among bottomonia




Study $e^+e^- \rightarrow \pi^+\pi^-\mu^+\mu^-$ (+ γ undetected)

- $Y(4S) \rightarrow \pi^+\pi^- Y(nS)$
- $e^+e^- \rightarrow \gamma_{\text{ISR}} Y(mS), Y(mS) \rightarrow \pi^+\pi^- Y(nS)$

Compare with Belle, 496 fb^{-1} [PRD 96 (2017) 5, 052005]

- Improved low momentum tracking



Junhao Yin, July 2021

Near term plan

By 2022 Belle II should have as much $Y(4S)$ as Belle. Analysis results from 2023!

Many analysis already ongoing, just waiting for more data!

Charmonium

- $X(3872)$ lineshape combining Belle and Belle II dataset
- Full amplitude analysis of $B \rightarrow$ charmonium modes
- Inclusive $B \rightarrow K(c\bar{c})$


Bottomonium sector

- Dalitz analysis of $Y(4S) \rightarrow \pi^+\pi^- Y(nS)$
- $h_b(1P)$ and $\eta_b(1S)$ exclusive decays

Junhao Yin, July 2021

Outside $Y(4S)$ even small data sets can make a difference

Currently available data sets in the bottomonium region




Junhao Yin, July 2021

JHEP10(2019)220 (Belle):

- “High-stat” scan points: 1 fb^{-1} each
- 1 point “on resonance”
- 2-3 points in the region of interest
- Significance: 5.2σ


Parameters:

	$\Upsilon(10860)$	$\Upsilon(11020)$	New structure
$M \text{ (MeV}/c^2)$	$10885.3 \pm 1.5^{+2.2}_{-0.9}$	$11000.0^{+4.0}_{-4.5} {}^{+1.0}_{-1.3}$	$10752.7 \pm 5.9^{+0.7}_{-1.1}$
$\Gamma \text{ (MeV)}$	$36.6^{+4.5}_{-3.9} {}^{+0.5}_{-1.1}$	$23.8^{+8.0}_{-6.8} {}^{+0.7}_{-1.8}$	$35.5^{+17.6}_{-11.3} {}^{+3.9}_{-3.3}$



Junhao Yin, July 2021

- Unlikely to be a molecule as it's far from any S- threshold
- No direct matching to conventional states (but may be an S-D mixing?)



Junhao Yin, July 2021

Summary

- We are at the beginning of a long program of quarkonium physics
 - Many opportunities for world-leading physics
 - Rediscoveries display the foundations we will build upon.
 - The legacy of Belle&Babar inspire us; LHCb&BESIII will push us as well.
 - 50 ab^{-1} integrated luminosity at Belle II in 6 years.
 - Expectation of great achievement in hadronic spectroscopy
 - Dedicated study of known XYZ states
 - Search for new particle via ISR, two photon production, double charmonium production...
 - Bottomonium search through $\Upsilon(nS)$
 - etc...

Junhao Yin, July 2021