



BELLE2-NOTE-PL-2020-023
DRAFT Version 1.0
July 28, 2020

Approved plots for $B \rightarrow \tau\nu$ study with 34.6 fb^{-1} of Phase III data

The Belle II Collaboration

Abstract

This note contains approved plots of $B \rightarrow \tau\nu_\tau$ analysis with ICHEP 2020 dataset, corresponding to an integrated luminosity of 34.6 fb^{-1} . Details of the analysis are documented in the supporting physics note BELLE2-NOTE-PH-2020-024.




FIG. 1: M_{bc} distribution of the tag B candidate for the electron channel. We require a reconstructed tag B with a FEI discriminant output $p_{FEI} > 0.01$ and only one charged track on the signal side. The track is required to have momentum in the lab frame $p_{trk} > 0.5 \text{ GeV}/c$ and to be identified as electron requiring $eid > 0.9$. A continuum suppression is applied requiring $\cos \Delta\theta_{\text{thrust}} < 0.8$. MC distributions are scaled to data luminosity.




FIG. 2: Distribution of the residual energy in the ECL, E_{ECL} , for the electron channel. E_{ECL} is the sum of the energy deposited in the ECL by all neutral objects in the event that are not used to make the tag B candidate or the signal side. Energy deposits greater than 55 MeV are considered. We require a reconstructed tag B with a FEI discriminant output $p_{FEI} > 0.01$, $M_{bc} > 5.27 \text{ GeV}/c^2$ and only one charged track on the signal side. The track is required to have momentum in the lab frame $p_{trk} > 0.5 \text{ GeV}/c$ and to be identified as electron requiring $eid > 0.9$. A continuum suppression is applied requiring $\cos \Delta\theta_{thrust} < 0.8$. MC distributions are scaled to data luminosity except for the signal which is enhanced by a factor of 100.




FIG. 3: Distribution of the sum of missing energy and missing momentum in the center-of-mass frame, $E_{\text{miss}}^* + cp_{\text{miss}}^*$, for the electron channel. We require a reconstructed tag B with a FEI discriminant output $p_{FEI} > 0.01$, $M_{bc} > 5.27 \text{ GeV}/c^2$ and only one charged track on the signal side. The track is required to have momentum in the lab frame $p_{trk} > 0.5 \text{ GeV}/c$ and to be identified as electron requiring $eid > 0.9$. A continuum suppression is applied requiring $\cos \Delta\theta_{\text{thrust}} < 0.8$. MC distributions are scaled to data luminosity except for the signal which is enhanced by a factor of 100.




FIG. 4: Monte Carlo distribution of the residual energy in the ECL, E_{ECL} , for the electron channel in the range $[0,2]$ GeV. E_{ECL} is the sum of the energy deposited in the ECL by all neutral objects in the event that are not used to make the tag B candidate or the signal side. Energy deposits greater than 55 MeV are considered. We require a reconstructed tag B with a FEI discriminant output $p_{FEI} > 0.01$, $M_{bc} > 5.27 \text{ GeV}/c^2$ and only one charged track on the signal side. The track is required to have momentum in the lab frame $p_{trk} > 0.5 \text{ GeV}/c$ and to be identified as electron requiring $eid > 0.9$. A continuum suppression is applied requiring $\cos \Delta\theta_{thrust} < 0.8$.